

First πK atom lifetime measurement and recent results from the DIRAC experiment

Monday, 2 June 2014 10:00 (0:30)

Collaboration

DIRAC

Abstract content

Low-energy QCD and specifically Chiral Perturbation Theory (ChPT) calculated $\pi\pi$ and πK scattering lengths with per cent precision. For processes involving u - and d -quarks theoretical predictions have been experimentally checked by $\pi^+\pi^-$ atom lifetime measurement [1] and by analysis of K -decays [2,3]. Detection and lifetime measurement of πK atom cast a look into processes which involve s -quark as well. We report evidence for πK atoms production, using 24 GeV/c proton beam from CERN PS interacting with a thin Ni target. We have identified (178 ± 49) πK pairs, which were produced and were subsequently broken-up (ionized) in the Ni target. Analysis yields a first measurement of the πK atom lifetime $(2.5_{-1.8}^{+3.0})$ fs [4]. This lifetime is connected in a model-independent way to the S-wave isospin-odd πK scattering lengths difference $|a_0^-| = \frac{1}{3}|a_{1/2} - a_{3/2}| = (0.11_{-0.04}^{+0.09}) M_\pi^{-1}$ (a_I for isospin I). Through the measurement of the $\pi^+\pi^-$ atom (pionium) lifetime, the experiment obtained the S-wave $\pi^+\pi^-$ scattering lengths difference $|a_0 - a_2|$ with 4% precision [1]. In 2011-2012 DIRAC collaboration collected data towards observation of long-lived (metastable) states of pionium. The observation of long-lived states opens the possibility to measure the energy difference between ns and np states and to determine the value of the combination $(2a_0 + a_2)$ of S-wave $\pi\pi$ scattering lengths. The experiment used two targets method: after production in the beryllium foil, atoms flew through a permanent magnetic field to reach the platinum ionization foil. The distance between foils is large enough for ns-states to vanish due to annihilation. Only $\pi^+\pi^-$ atoms in states with non-zero angular momentum can get into the second Pt target. We report unambiguous observation of long-lived (metastable) states of $\pi^+\pi^-$ atoms.

[1] B. Adeva, et al. (DIRAC Collaboration), Phys. Lett. B704 (2011) 24. [2] J.R. Batley, et al. (NA48/2 Collaboration), Eur. Phys. J. C64 (2009) 589. [3] J.R. Batley, et al. (NA48/2 Collaboration), Eur. Phys. J. C70 (2010) 635. [4] B. Adeva, et al. (DIRAC Collaboration), arXiv:1403.0845.

Primary author(s) : ZHABITSKY, Mikhail (Joint Institute for Nuclear Research)

Presenter(s) : ZHABITSKY, Mikhail (Joint Institute for Nuclear Research)

Session Classification : Plenary Session