

Predictions on the second-class current decays

$$\tau^- \rightarrow \pi^- \eta^{(\prime)} \nu_\tau$$

Saturday, 4 June 2016 16:00 (1:30)

Collaboration

Abstract content

We analyze the second-class current decays $\tau^- \rightarrow \pi^- \eta^{(\prime)} \nu_\tau$ in the framework of Chiral Perturbation Theory with Resonances. Taking into account $\pi^0 - \eta - \eta'$ mixing, the $\pi - \eta^{(\prime)}$ vector form factor is extracted, in a model-independent way, using existing data on the $\pi - \pi_0$ one. For the participant scalar form factor, we have considered different parameterizations ordered according to their increasing fulfillment of analyticity and unitarity constraints. We start with a Breit-Wigner parameterization dominated by the $a_0(980)$ scalar resonance and after we include its excited state, the $a_0(1450)$. We follow by an elastic dispersion relation representation through the Omnes integral. Then, we illustrate a method to derive a closed-form expression for the $\pi - \eta$, $\pi - \eta'$ (and $K - K^0$) scalar form factors in a coupled-channels treatment. Finally, predictions for the branching ratios and spectra are discussed emphasizing the error analysis. An interesting result of this study is that both $\tau^- \rightarrow \pi^- \eta^{(\prime)} \nu_\tau$ decay channels are promising for the soon discovery of second-class currents at Belle-II. We also predict the relevant observables for the partner $\eta^{(\prime)}_{13}$ decays, which are extremely suppressed in the Standard Model.

Primary author(s) : ESCRIBANO, Rafel (Universitat Autònoma de Barcelona)

Co-author(s) : GONZÁLEZ-SOLÍS, Sergi (Universitat Autònoma de Barcelona); ROIG, Pablo (CINVESTAV (México))

Presenter(s) : GONZÁLEZ-SOLÍS, Sergi (Institut de Física d'Altes Energies)

Session Classification : Poster Session