

Inclusive production of J/Ψ and Ψ' mesons at the LHC

Anna Cisek

University of Rzeszow

MESON 2016, Krakow, 2 - 7 June 2016

Outline

1 Introduction

2 Formalism

- diagrams
- J/Ψ and Ψ' mesons production
- J/Ψ production from radiative decay of χ_c mesons

3 Results

- J/Ψ
- Ψ'
- χ_c


4 Conclusions

Introduction


- There is a long-standing lack of convergence in understanding production of J/ψ quarkonia in proton-proton or proton-antiproton collisions
- Some authors believe that the corresponding cross sections **are dominated by the so-called color-octet contribution**
- Some other authors expect that the **color-singlet contribution dominates**
- The **color-octet contribution** cannot be calculated from first principle and **is rather fitted to the experimental data**
- **We calculate the color-singlet contribution** as well as possible in the NRQCD k_t -factorization
- We concentrate rather on small transverse momenta of J/ψ or ψ' relevant for ALICE and LHCb data

The possible mechanism to production of vector meson in hadronic collisions (exclusive processes)


Photoproduction



Oderon-Pomeron fusion



Radiative Decay of χ_c




Khoze-Martin-Ryskin 2002
 Klein-Nystrand 2004
 Schäfer, Szczurek 2007
 (W. Schäfer talk)

Schäfer, Mankiewicz, Nachtmann 1991
 Bzdak, Motyka, Szymanowski, Cudell 2007

Pasechnik, Szczurek, Teryaev 2008

The main color-singlet mechanism of production of J/Ψ and Ψ' mesons



- We restrict to gluon-gluon fusion mechanism

Differential cross section for J/Ψ


- In the NLO differential cross section in the k_t factorization can be written as:

$$\frac{d\sigma(pp \rightarrow J/\psi g X)}{dy_{J/\psi} dy_g d^2 p_{J/\psi,t} d^2 p_{g,t}} = \frac{1}{16\pi^2 \hat{s}^2} \int \frac{d^2 q_{1t}}{\pi} \frac{d^2 q_{2t}}{\pi} \overline{|\mathcal{M}_{\mathbf{g}^* \mathbf{g}^* \rightarrow \mathbf{J}/\psi \mathbf{g}}|^2} \times \\ \times \delta^2 (\mathbf{q}_{1t} + \mathbf{q}_{2t} - \mathbf{p}_{\mathbf{J}/\psi, t} - \mathbf{p}_{\mathbf{g}, t}) \mathcal{F}_g(\mathbf{x}_1, \mathbf{q}_{1t}^2, \mu_F^2) \mathcal{F}_g(\mathbf{x}_2, \mathbf{q}_{2t}^2, \mu_F^2)$$

- We calculate the dominant color-single $gg \rightarrow J/\psi g$ contribution taking into account transverse momenta of initial gluons
- The corresponding matrix element squared for the $gg \rightarrow J/\psi g$ is

$$|\mathcal{M}_{gg \rightarrow J/\psi g}|^2 \propto \alpha_s^3 |\mathbf{R}(\mathbf{0})|^2$$

Matrix elements for J/Ψ



$$\begin{aligned} \mathcal{M}_a(\text{gg} \rightarrow J/\psi g) = & tr\{\epsilon_1(\mathbf{p}_c - \mathbf{k}_1 + m_c)\epsilon_2 \times (-\mathbf{p}_c - \mathbf{k}_3 + m_c)\epsilon_3 J(S, L)\} C_\Psi \\ & \times tr\{T^a T^b T^c T^d\} [k_1^2 - 2(p_c k_1)]^{-1} \times [k_3^2 - 2(p_{\bar{c}} k_3)]^{-1} + 5 \text{ permutations} \end{aligned}$$

S. P. Baranov, Phys. Rev. D **66** (2002) 114003

Matrix elements for J/Ψ

$$\begin{aligned}\mathcal{M}_b(\text{gg} \rightarrow J/\psi \text{g}) = & \text{tr}\{\gamma_\mu(p_{\bar{c}} - k_3 + m_c)\epsilon_3 J(S, L)\} \\ & \times G^3(k_1, \epsilon_1, k_2, \epsilon_2, -k, \mu) C_\Psi f^{abe} \\ & \times \text{tr}\{T^e T^c T^d\} [k^2]^{-1} \\ & \times [k_3^2 - 2(p_{\bar{c}} k_3)]^{-1} + 5 \text{ permutations}\end{aligned}$$

$$\begin{aligned}\mathcal{M}_c(\text{gg} \rightarrow J/\psi \text{g}) = & \text{tr}\{\gamma_\mu J(S, L)\} G^3(k_1, \epsilon_1, k_2, \epsilon_2, -k, \mu) \\ & \times G^3(-k_3, -\epsilon_3, -p_\Psi, -\epsilon_-, -k, \nu) C_\Psi f^{abe} f^{cfe} \\ & \times \text{tr}\{T^f T^d\} [k^2]^{-1} \times [m_\Psi^2]^{-1} + 2 \text{ permutations}\end{aligned}$$

$$\begin{aligned}\mathcal{M}_d(\text{gg} \rightarrow J/\psi \text{g}) = & \text{tr}\{\gamma_\nu J(S, L)\} G^{(4)A,B,C}(\epsilon_1, \epsilon_2, \epsilon_3, \nu) C_\Psi \\ & \times \text{tr}\{T^f T^d\} [k^2]^{-1} [m_\Psi^2]^{-1}\end{aligned}$$

S. P. Baranov, Phys. Rev. D **66** (2002) 114003

χ_c production

- * In the k_t -factorization approach the leading-order **cross section for the χ_c meson production** can be written as:

$$\sigma_{\text{pp} \rightarrow \chi_c} = \int \frac{dx_1}{x_1} \frac{dx_2}{x_2} \frac{d^2 q_{1t}}{\pi} \frac{d^2 q_{2t}}{\pi} \delta((q_1 + q_2)^2 - M_{\chi_c}^2) \sigma_{gg \rightarrow H}(x_1, x_2, q_1, q_2) \\ \times \mathcal{F}_g(\mathbf{x}_1, \mathbf{q}_{1t}^2, \mu_F^2) \mathcal{F}_g(\mathbf{x}_2, \mathbf{q}_{2t}^2, \mu_F^2)$$

- * The matrix element squared for the $gg \rightarrow \chi_c$ subprocess is

$$|\mathcal{M}_{gg \rightarrow \chi_c}|^2 \propto \alpha_s^2 |\mathbf{R}'(\mathbf{0})|^2$$

- * For running coupling constants we choose:

$$\alpha_s^2 \rightarrow \alpha_s(\mu_1^2) \alpha_s(\mu_2^2)$$

where $\mu_1^2 = \max(\mathbf{q}_{1t}^2, \mathbf{m}_t^2)$ and $\mu_2^2 = \max(\mathbf{q}_{2t}^2, \mathbf{m}_t^2)$

Cross section for χ_c

- After some manipulation:

$$\sigma_{\text{pp} \rightarrow \chi_c} = \int dy d^2 p_t d^2 q_t \frac{1}{s \mathbf{x}_1 \mathbf{x}_2} \frac{1}{m_{t,\chi_c}^2} \overline{|\mathcal{M}_{\mathbf{g}^* \mathbf{g}^* \rightarrow \chi_c}|^2} \mathcal{F}_{\mathbf{g}}(\mathbf{x}_1, \mathbf{q}_{1t}^2, \mu_F^2) \mathcal{F}_{\mathbf{g}}(\mathbf{x}_2, \mathbf{q}_{2t}^2, \mu_F^2) / 4$$

- Which can be also used to calculate rapidity and transverse momentum distribution of the χ_c mesons
- In the last equation:

$$\mathbf{p}_t = \mathbf{q}_{1t} + \mathbf{q}_{2t} \quad \mathbf{q}_t = \mathbf{q}_{1t} - \mathbf{q}_{2t}$$

$$\mathbf{x}_1 = \frac{\mathbf{m}_{t,\chi_c}}{\sqrt{s}} \exp(\mathbf{y}) \quad \mathbf{x}_2 = \frac{\mathbf{m}_{t,\chi_c}}{\sqrt{s}} \exp(-\mathbf{y})$$

- The factor $\frac{1}{4}$ is the jacobian of transformation from $(\mathbf{q}_{1t}, \mathbf{q}_{2t})$ to $(\mathbf{p}_t, \mathbf{q}_t)$ variables

Matrix elements for χ_c

$$\begin{aligned}\overline{|\mathcal{A}(R + R \rightarrow \mathcal{H}[{}^3P_0^{(1)}])|^2} &= \frac{8}{3}\pi^2\alpha_s^2 \frac{\langle \mathcal{O}^{\mathcal{H}}[{}^3\mathbf{P}_0^{(1)}] \rangle}{M^5} \mathbf{F}^{{}^3\mathbf{P}_0}(\mathbf{t}_1, \mathbf{t}_2, \varphi) \\ \overline{|\mathcal{A}(R + R \rightarrow \mathcal{H}[{}^3P_1^{(1)}])|^2} &= \frac{16}{3}\pi^2\alpha_s^2 \frac{\langle \mathcal{O}^{\mathcal{H}}[{}^3\mathbf{P}_1^{(1)}] \rangle}{M^5} \mathbf{F}^{{}^3\mathbf{P}_1}(\mathbf{t}_1, \mathbf{t}_2, \varphi) \\ \overline{|\mathcal{A}(R + R \rightarrow \mathcal{H}[{}^3P_2^{(1)}])|^2} &= \frac{32}{45}\pi^2\alpha_s^2 \frac{\langle \mathcal{O}^{\mathcal{H}}[{}^3\mathbf{P}_2^{(1)}] \rangle}{M^5} \mathbf{F}^{{}^3\mathbf{P}_2}(\mathbf{t}_1, \mathbf{t}_2, \varphi)\end{aligned}$$

where

$$\langle \mathcal{O}^{\chi_{cJ}}[{}^3\mathbf{P}_{\mathbf{J}}^{(1)}] \rangle = 2N_c(2J+1)|\mathbf{R}'(\mathbf{0})|^2$$

B. A. Kniehl, D. V. Vasin, V. A. Saleev; Phys. Rev. D **73** (2006) 074022

Matrix elements for χ_c

$$\mathbf{F}^{[{}^3\mathbf{P}_0]}(\mathbf{t}_1, \mathbf{t}_2, \varphi) = \frac{2}{9} \frac{M^2 (M^2 + |\mathbf{p}_t|^2)^2 [(3M^2 + t_1 + t_2) \cos \varphi + 2\sqrt{t_1 t_2}]^2}{(M^2 + t_1 + t_2)^4}$$

$$\mathbf{F}^{[{}^3\mathbf{P}_1]}(\mathbf{t}_1, \mathbf{t}_2, \varphi) = \frac{2}{9} \frac{M^2 (M^2 + |\mathbf{p}_t|^2)^2 [(t_1 + t_2)^2 \sin^2 \varphi + M^2 (t_1 + t_2 - 2\sqrt{t_1 t_2} \cos \varphi)]}{(M^2 + t_1 + t_2)^4}$$

$$\begin{aligned} \mathbf{F}^{[{}^3\mathbf{P}_2]}(\mathbf{t}_1, \mathbf{t}_2, \varphi) = \frac{1}{3} \frac{M^2}{(M^2 + t_1 + t_2)^4} & (M^2 + |\mathbf{p}_t|^2)^2 \{ 3M^4 + 3M^2(t_1 + t_2) + 4t_1 t_2 \\ & + (t_1 + t_2)^2 \cos^2 \varphi + 2\sqrt{t_1 t_2} [3M^2 + 2(t_1 + t_2)] \cos \varphi \} \end{aligned}$$

where $\mathbf{p}_t = \mathbf{q}_{1t} + \mathbf{q}_{2t}$






and $\varphi = \varphi_1 - \varphi_2$ is the angle between \mathbf{q}_{1t} and \mathbf{q}_{2t} so

$$|\mathbf{p}_t|^2 = t_1 + t_2 + 2\sqrt{t_1 t_2} \cos \varphi$$


B. A. Kniehl, D. V. Vasin, V. A. Saleev; Phys. Rev. D **73** (2006) 074022

rapidity dependence for J/Ψ meson

- **2.76 TeV** - B. Abelev et al.;
Phys. Let. B. **718** (2012) 295-306
- **7 TeV** - B. Abelev et al.;
Eur.Phys. J. C. **74** (2014) 2974
7 TeV - K. Aamodt et al.;
Phys. Let. B. **704** (2011) 442
7 TeV - R. Aaij et al.;
Eur.Phys. J. C. **71** (2011) 1645
- **13 TeV** - R. Aaij et al.;
JHEP 1510 (2015) 172





p_t distribution for J/Ψ meson





- 7 TeV - R. Aaij et al.; Eur.Phys. J. C. 71 (2011) 1645

rapidity distribution for Ψ' meson




- **7 TeV** - B. Abelev et al.;
Eur.Phys. J. C. **74** (2014) 2974




- KMR alone overshoot experimental data for rapidity distribution
- the best solution is takes KMR distribution for large x and KS for small x

different UGDF for χ_c meson rapidity distribution



mixed UGDF for χ_c meson rapidity distribution



Conclusions

- We calculate the color-singlet contribution in the NRQCD k_t -factorization
- We compare our results with ALICE and LHCb data
- Our results in rapidity are almost consistent with experimental data
- Cross section strongly depends on UGDF function
- The best solution is to calculate mixed UGDFs (KMR-KS)
- Only small room is left for color-octet contribution