


Electromagnetic transition form factor of the η meson with WASA-at-COSY



Ankita Goswami

(for the WASA-at-COSY collaboration)

Indian Institute of Technology Indore



MESON 2016

14th International Workshop on Meson Production,
Properties and Interaction

Motivation


Intrinsic structure of hadrons

form factors


Vector meson dominance

background for physics beyond standard model

rare pion decay $\pi^0 \rightarrow e^+ e^-$



g-2 of muon





Transition Form Factor

Transition Form Factor $F(q^2)$ of the η meson is observed through the rare electromagnetic decay $\eta \rightarrow \gamma e^+ e^-$ (BR $\rightarrow 6.9 \times 10^{-3}$).

$$\frac{d\Gamma(\eta \rightarrow \gamma e^+ e^-)}{dq^2 \cdot \Gamma(\eta \rightarrow \gamma\gamma)} = \frac{2\alpha}{3\pi} \left[1 - \frac{4m_e^2}{q^2}\right]^{1/2} \left[1 + \frac{2m_e^2}{q^2}\right] \frac{1}{q^2} \left[1 - \frac{q^2}{m_\eta^2}\right]^3 |F_\eta(q^2)|^2$$

$$F(q^2) = \frac{1}{1 - \frac{q^2}{\Lambda^2}} \approx 1 + \frac{q^2}{\Lambda^2} \quad \left| \frac{dF(q^2)}{dq^2} \right|_{q^2=0} = \frac{1}{\Lambda^2} = b_\eta$$

Λ is pole mass and b_η is slope of the form factor




PHYSICAL REVIEW C 89, 044608 (2014)

$$\Lambda^2 = (1.95 \pm 0.15_{\text{stat}} \pm 0.10_{\text{syst}}) \text{ GeV}^{-2}$$

WASA (Wide Angle Shower Apparatus) set up



Reaction: $p + p \rightarrow p + p + \eta(e^+ e^- \gamma)$ at beam energy 1.4 GeV



- Fixed target experiment, pellet target, 22.9 % of 4π acceptance
- Recoil protons are detected with the forward detector
- e^+e^- are detected with the mini drift chamber in the magnetic field of solenoid
- Photons are detected in the calorimeter

Data Analysis: Particle Identification


$$p + p \rightarrow p + p + \eta(e^+ e^- \gamma)$$



- Protons are identified in the forward part of the detector
- Deposit energy in forward range hodoscope layers

- Different types of particles leave distinct bands
- Momentum times charge of the particle is plotted against the energy deposited by particle in the calorimeter

Energy-momentum balance





Missing Energy:

$$E_{\text{target}} + E_{\text{beam}} - (E_{\text{proton1}} + E_{\text{proton2}} + E_{e^+} + E_{e^-} + E_{\gamma})$$





Missing Momentum:

$$\mathbf{P}_{\text{target}} + \mathbf{P}_{\text{beam}} - (\mathbf{P}_{\text{proton1}} + \mathbf{P}_{\text{proton2}} + \mathbf{P}_{e^+} + \mathbf{P}_{e^-} + \mathbf{P}_{\gamma})$$

Background suppression:
event candidates will still have pions





Conversion background



- Photons interact with beam-pipe material and convert into $e^+ e^-$ pairs
- $\eta \rightarrow \gamma\gamma$ contributes
- Invariant mass at beam pipe plotted against the radius of closest approach of $e^+ e^-$



Split off background



- Photons and electrons make electromagnetic shower in the calorimeter
- Split-offs are discontinuous showers
- We look at the energy deposited in the calorimeter v/s the angle between photon candidate and closest charged track

split offs are located at low energy and small angle


Missing mass of η meson



- Main background source is $pp \rightarrow pp\pi^0\pi^0$ (π^0 Dalitz decay)
- Background fit: $\text{pol4} \times \text{MC}$ ($pp \rightarrow pp\pi^0\pi^0$ (π^0 Dalitz decay)) excluding the peak region
- produced η : 10^8
- approximately 43k η decays

Background study: cocktail plots

preliminary and not acceptance corrected




Background channel	Cross-section/ Branching ratio	Probability of being detected as signal (%)
$pp \rightarrow pp\pi^0(e^+e^-\pi^0)(\gamma\gamma)$	324 μb	.069
$pp \rightarrow pp\pi^+\pi^-\pi^0(e^+e^-\gamma)$	4.6 μb	.00041
$pp \rightarrow pp\pi^0(e^+e^-\gamma)\pi^0(\gamma\gamma)$	1.34 μb	.011
$\eta \rightarrow \pi^+\pi^-\pi^0$	22.6 %	.0009
$\eta \rightarrow \pi^+\pi^-\gamma$	4.68 %	.0287
$\eta \rightarrow \gamma\gamma$	39 %	.0032
$\eta \rightarrow \pi^0(\gamma\gamma)\pi^0(\gamma\gamma)\pi^0(e^+e^-\gamma)$	32 %	.122

- *Direct and competing decays*
- *Phase space simulations (for now)*
- *Δ - Δ , $\pi^+\pi^-\pi^0$ correlations have to be implemented*
- *Normalization of background channels is done relative to each other and scaled with data*

reaching for the double Dalitz decay

Susan Schadmand

pp η 2010 | $\eta \rightarrow e^+ e^- e^+ e^-$ | cut-based analysis: background study



- WASA-at-COSY standard analysis
- preliminary and not acceptance corrected.
- **consistency-check** : yield consistent with our preliminary single Dalitz decay analysis

goal: evaluate branching ratio

latest WASA result: nucl-ex/1509.06588
 $BR = (3.2 \pm 0.9_{stat} \pm 0.5_{sys}) \times 10^{-5}$

Summary

$$\eta \rightarrow \gamma e^+ e^-$$

- Main source of background is $pp \rightarrow pp \pi^0 \pi^0 (\pi^0 \rightarrow e^+ e^- \gamma)$
- Detailed study of background channels is ongoing

$$\eta \rightarrow e^+ e^- e^+ e^-$$

- Branching ratio

Outlook

- As a different approach, kinematic fit to suppress background
- Transition form factor of η