## Partial wave analysis of $\pi\pi$ scattering below 2 GeV

Friday, 3 June 2016 18:00 (0:20)

## Collaboration

## Abstract content

In a unitary multi-channel approach, precise determination of  $\pi\pi$  scattering amplitudes for D and F waves has been presented. These scattering amplitudes are in the  $I^G J^{PC} = 0^+ 2^{++}$  sector on the processes of  $\pi\pi \to \pi\pi$ ,  $4\pi$ ,  $K\bar{K}$  and  $\eta\eta$ , likewise in the  $I^G J^{PC} = 1^+ 3^{--}$  sector on the processes of  $\pi\pi \to \pi\pi$ ,  $4\pi$ ,  $\omega\pi$  and  $K\bar{K}$  for D and F waves respectively. The amplitudes were refined and re-fitted to the dispersion relations up to 1.1 GeV, and to the experimental data in the effective two pion mass from the threshold to 2.7 GeV and 1.9 GeV for D and F waves, respectively. Old parameterizations did not satisfy the crossing symmetry condition and did not describe the  $\pi\pi$  threshold region. Moreover, a satisfactory justification regarding the controversies in the states of  $f_2$  and  $\rho_3$  mesons about their masses and number of states that are taken into account has been discussed and finalized.

Primary author(s) :NAZARI, Vahabeddin (INP PAN Cracow)Presenter(s) :NAZARI, Vahabeddin (INP PAN Cracow)Session Classification :Parallel Session B4