

$a_0(980)$ photoproduction in the coupled channel model

Friday, 3 June 2016 17:00 (0:20)

Collaboration

Abstract content

We constructed the $\gamma p \rightarrow p\pi\eta$ S-wave photoproduction amplitudes which, for the first time in the literature, take into account effects of the $K\bar{K}-\pi\eta$ interchannel coupling in effective mass region corresponding to the $a_0(980)$ resonance [1]. The principal merit of our approach is that apart from the coupling constants already successfully applied in other analyses [2] and resonance parameters like masses, widths and branching ratios used in the final state interaction parametrisation [3] we do not engage any new model parameters.

We also calculated the Born amplitudes and cross sections for partial waves P and D. These can give an estimation of the photoproduction cross section of spin 1 and spin 2 $\pi\eta$ resonances like $\pi_1(1400)$ and $a_2(1320)$.

It is important to stress that reliable model of the photoproduction in the $\pi\eta$ channel is essential for proper description of data to be collected by new JLab experiments CLAS12 and GlueX.

Literature:

- [1] Ł. Bibrzycki, R. Kamiński, <http://arxiv.org/abs/1509.06135>
- [2] Ł. Bibrzycki and R. Kamiński, Phys. Rev. D87, 114010 (2013)
- [3] A. Furman, L. Leśniak, Phys.Lett. B538, 266 (2002)

Primary author(s) : BIBRZYCKI, Łukasz (Pedagogical University of Cracow); KAMIŃSKI, Robert (Institute of Nuclear Physics PAN, Kraków)

Presenter(s) : BIBRZYCKI, Łukasz (Pedagogical University of Cracow)

Session Classification : Parallel Session B4