

Charm physics prospects at the Belle II experiment

Thursday, 7 June 2018 18:15 (0:20)

Collaboration

Belle II

Abstract content

The Belle II experiment is a substantial upgrade of the Belle detector and will operate at the SuperKEKB energy-asymmetric e^+e^- collider. The accelerator has already successfully completed the first phase of commissioning in 2016 and first electron-positron collisions in Belle II are expected for April 2018. The design luminosity of SuperKEKB is $8 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ and the Belle II experiment aims to record 50 ab^{-1} of data, a factor of 50 more than the Belle experiment. In this presentation, we will discuss the expected sensitivity of Belle II for CPV measurements and New Physics searches in the charm sector. Alternative flavor-tagging techniques have been developed, a novel flavor-tagging method of prompt D^0 s will be presented. Finally, we will present the impact of the improved tracking at Belle II, that will allow to significantly increase the precision of time-dependent measurements. This abstract is presented on behalf of the Belle II collaboration; if accepted the Speakers Committee will assign the talk to a speaker.

Primary author(s) : STARIC, Marko (J. Stefan Institute)

Presenter(s) : STARIC, Marko (J. Stefan Institute)

Session Classification : Parallel Session A2