

# NEW INSIGHTS ON LOW ENERGY $\pi N$ SCATTERING AMPLITUDES

**Meson 2018 @ Kraków**

Y.F. Wang

based on **arXiv:1712.09257**, in collaboration with **D.L. Yao** and **H.Q. Zheng**

School of physics, Peking University

June 7, 2018



# CONTENTS

- 1 Introduction
  - The pion-nucleon scattering
  - PKU representation
- 2 Theoretical framework
  - $\pi N$  partial wave amplitudes
  - Phase shift components
- 3 Numerical results
  - Tree-level qualitative analyses
  - Discrepancies in  $S_{11}$  and  $P_{11}$  channels
- 4 Hidden contributions
  - $P_{11}$  channel: shadow pole of the nucleon
  - $S_{11}$  channel: lowest potential-nature resonance?
- 5 Summary





# 1. Introduction

# THE PION-NUCLEON SCATTERING





- The  $\pi N$  scattering  $\rightarrow$  one of the most fundamental and important processes in nuclear or hadron physics
- Decades of researching
- Various experiments and phenomena  
( $L_{2I} 2J$  convention,  $W = \sqrt{s}$ ,  $S_r = 1 - \eta^2$ )[SAID: WI 08]





# THE PION-NUCLEON SCATTERING



# THE PION-NUCLEON SCATTERING



# THE PION-NUCLEON SCATTERING



# THEORETICAL DISCUSSIONS

- Problems to study

- Low energy properties:

- $\pi N$   $\sigma$ -term, subthreshold expansions

- [C. Ditsche et. al. 2012 JHEP][Y. H. Chen et. al. 2013 PRD][Hoferichter et. al. 2016 Phys.Rept.]

- Intermediate resonances:  $\Delta(1232)$ ,  $N^*(1440)$ ,  $N^*(1535) \dots$

- Methods

- Perturbative calculation

- [J.M. Alarcón et. al. 2012 RPD][Y. H. Chen et. al. 2013 PRD]

- Couple channel Lippmann-Schwinger Equation

- [O. Krehl et. al. 2000 PRC]

- Dispersion technique [A. Gasparyan and M.F.M. Lutz 2010 NPA]

- Roy-Steiner equation

- [C. Ditsche et. al. 2012 JHEP][Hoferichter et. al. 2016 Phys.Rept.]



# $S_{11}$ AND $P_{11}$ CHANNELS

- $S_{11}$  channel ( $L_{2I} 2J$  convention):  $N^*(1535)$   
[N. Kaiser et. al. 1995 PLB][J. Nieves et. al. 2000 PRD]
  - lies above the  $P$ - wave first resonance  $N^*(1440)$
  - large couple channel effects with  $\pi N$  and  $\eta N$
- $P_{11}$  channel:  $N^*(1440)$  (Ropper resonance), various puzzles
  - low mass, large decay width, coupling to  $\sigma N$  channel...  
[O. Krehl et. al. 2000 PRC]
  - two-pole structure? [R. A. Arndt et. al. 1985 PRD]
  - second sheet complex branch cut in  $P_{11}$  channel?  
[S. Ceci et. al. 2011 PRC]
- A method is needed to examine the relevant channels carefully and to exhume more physics behind
  - low energy
  - model independent

# PKU REPRESENTATION

- Peking University (PKU) representation: elastic two-body scatterings

$$S = \prod_i S_i \times S_{cut}$$

- $S_i$ : pole terms,  $S_{cut} = e^{2i\rho(s)f(s)}$ : left-hand cuts and right hand inelastic cut – background.

$$f(s) = \frac{s}{2\pi i} \int_L ds' \frac{\text{disc}f(s')}{(s' - s)s'} + \frac{s}{2\pi i} \int_{R'} ds' \frac{\text{disc}f(s')}{(s' - s)s'}$$

- $f(0) \equiv 0$  [Z. Y. Zhou and H. Q. Zheng 2006 NPA]

# PKU REPRESENTATION

- $f(s)$  perturbatively calculated, poles as parameters (input or fit)
- Corresponding to the Ning Hu representation in QM

[N. Hu 1948 PR]

- Advantages

- rigorous and universal
- separated  $S \rightarrow$  additive phase shift
- sensitive to (not too) distant poles
- definite sign of the phase shifts  $\rightarrow$  figuring out hidden contributions

- Applications


- the  $\pi\pi$  elastic scattering  $\rightarrow$  existence of the  $\sigma$  particle ( $f_0(500)$ ) [Z. G. Xiao and H. Q. Zheng 2001 NPA]
- the  $\pi K$  elastic scattering  $\rightarrow$   $\kappa$  resonance ( $K^*(800)$ )

[H. Q. Zheng et. al. 2004 NPA]

# THE EXISTENCE OF $\sigma$

The left-hand cut contribution (negative definite)

→ the existence of  $\sigma$  particle [Z. G. Xiao and H. Q. Zheng 2001 NPA]



$$M_\sigma = 457 \pm 15 \text{ MeV}, \Gamma_\sigma = 551 \pm 28 \text{ MeV} \quad [\text{Z. Y. Zhou et. al 2005 JHEP}]$$

$$M_\sigma = 441_{-8}^{+16} \text{ MeV}, \Gamma_\sigma = 544_{-25}^{+18} \text{ MeV} \quad [\text{I. Caprini et. al. 2006 PRL}]$$

## 2. Theoretical framework

# LAGRANGIAN

- Covariant baryon chiral perturbation theory,  $SU(2)$  case.
- Lagrangians [N. Fettes et. al. 2000 Ann. Phys. ]
- $\mathcal{O}(p^1)$ :

$$\mathcal{L}_{\pi N}^{(1)} = \bar{N}(i\not{D} - M + \frac{1}{2}g\not{u}\gamma^5)N$$

- $\mathcal{O}(p^2)$  (“ $\langle \rangle$ ” stands for trace in isospin space):

$$\begin{aligned}\mathcal{L}_{\pi N}^{(2)} = & c_1 \langle \chi_+ \rangle \bar{N}N - \frac{c_2}{4M_N^2} \langle u^\mu u^\nu \rangle (\bar{N}D_\mu D_\nu N + \text{h.c.}) \\ & + \frac{c_3}{2} \langle u^\mu u_\mu \rangle \bar{N}N - \frac{c_4}{4} \bar{N} \gamma^\mu \gamma^\nu [u^\mu, u^\nu] N\end{aligned}$$

# CONVENTIONS

- Conventions

$$D_\mu = \partial_\mu + \Gamma_\mu$$

$$\Gamma_\mu = \frac{1}{2} [u^\dagger (\partial_\mu - ir_\mu) u + u (\partial_\mu - il_\mu) u^\dagger]$$

$$u_\mu = i [u^\dagger (\partial_\mu - ir_\mu) u - u (\partial_\mu - il_\mu) u^\dagger]$$

$$\chi_\pm \equiv u^\dagger \chi u^\dagger \pm u \chi^\dagger u$$

$$\chi = 2B_0(s + ip)$$

$$h_\nu^\mu = [D_\nu, u^\mu] + [D^\mu, u_\nu]$$

In calculation  $2B_0s \rightarrow 2B_0m_q = m_\pi^2$ , other sources are switched off

# ISOSPIN DECOMPOSITION

- Symmetric part vs. anti-symmetric part

$$T(\pi^a + N_i \rightarrow \pi^{a'} + N_f) = \chi_f^\dagger \left( \delta^{aa'} T^S + \frac{1}{2} [\tau^{a'}, \tau^a] T^A \right) \chi_i$$

- Isospin channels

$$T^{I=1/2} = T^S + 2T^A$$

$$T^{I=3/2} = T^S - T^A$$



# HELICITY STRUCTURE

- Lorentz structure

$$\begin{aligned}T^{S,A} &= \bar{u}(p', s') \left[ A^{S,A}(s, t) + \frac{1}{2}(\not{q} + \not{q}') B^{S,A}(s, t) \right] u(p, s) \\ &= \bar{u}(p', s') \left[ D^{S,A}(s, t) + \frac{i\sigma^{\mu\nu} q_\nu q'_\mu}{2M} B^{S,A}(s, t) \right] u(p, s)\end{aligned}$$

where  $D = A + (s - u)B/(4M_N)$

- Helicity amplitudes ( $z_s = \cos \theta$ )

$$T_{++} = \left(\frac{1+z_s}{2}\right)^{\frac{1}{2}} [2M_N A(s, t) + (s - m_\pi^2 - M_N^2) B(s, t)]$$

$$T_{+-} = -\left(\frac{1-z_s}{2}\right)^{\frac{1}{2}} s^{-\frac{1}{2}} [(s - m_\pi^2 + M_N^2) A(s, t) + M_N (s + m_\pi^2 - M_N^2) B(s, t)]$$

- Partial wave projection

$$T_{++}^J = \frac{1}{32\pi} \int_{-1}^1 dz_s T_{++}(s, t(s, z_s)) d_{-1/2, -1/2}^J(z_s)$$

$$T_{+-}^J = \frac{1}{32\pi} \int_{-1}^1 dz_s T_{+-}(s, t(s, z_s)) d_{1/2, -1/2}^J(z_s)$$

# CHANNELS TO BE ANALYZED

$L_{2I\ 2J}$  convention

$$T(S_{11}) = T_{++}(I = 1/2, J = 1/2) + T_{+-}(I = 1/2, J = 1/2)$$

$$T(S_{31}) = T_{++}(I = 3/2, J = 1/2) + T_{+-}(I = 3/2, J = 1/2)$$

$$T(P_{11}) = T_{++}(I = 1/2, J = 1/2) - T_{+-}(I = 1/2, J = 1/2)$$

$$T(P_{31}) = T_{++}(I = 3/2, J = 1/2) - T_{+-}(I = 3/2, J = 1/2)$$

$$T(P_{13}) = T_{++}(I = 1/2, J = 3/2) + T_{+-}(I = 1/2, J = 3/2)$$

$$T(P_{33}) = T_{++}(I = 3/2, J = 3/2) + T_{+-}(I = 3/2, J = 3/2)$$

Each channel satisfies unitarity condition.

# PKU REPRESENTATION

- PKU representation

$$S(s) = \prod_b \frac{1 - i\rho(s) \frac{s}{s-s_L} \sqrt{\frac{s_b-s_L}{s_R-s_b}}}{1 + i\rho(s) \frac{s}{s-s_L} \sqrt{\frac{s_b-s_L}{s_R-s_b}}} \prod_v \frac{1 + i\rho(s) \frac{s}{s-s_L} \sqrt{\frac{s'_v-s_L}{s_R-s'_v}}}{1 - i\rho(s) \frac{s}{s-s_L} \sqrt{\frac{s'_v-s_L}{s_R-s'_v}}} \\ \prod_r \frac{M_r^2 - s + i\rho(s)sG_r}{M_r^2 - s - i\rho(s)sG_r} e^{2i\rho(s)f(s)}$$

- $s_b$ : bound states.  $s'_v$ : virtual states (sheet I).  $z_r$ : resonances (sheet II).
- $s_L = (m_1 - m_2)^2$ ,  $s_R = (m_1 + m_2)^2$ ,  $\rho(s) = \sqrt{s - s_L} \sqrt{s - s_R} / s$ .

$$M_r^2 = \operatorname{Re}[z_r] + \operatorname{Im}[z_r] \frac{\operatorname{Im}[\sqrt{(z_r - s_R)(z_r - s_L)}]}{\operatorname{Re}[\sqrt{(z_r - s_R)(z_r - s_L)}]} \\ G_r = \frac{\operatorname{Im}[z_r]}{\operatorname{Re}[\sqrt{(z_r - s_R)(z_r - s_L)}]}$$


# PHASE SHIFT COMPONENTS

- PKU representation  $\rightarrow$  conventionally additive phase shift
- Phase shift contributions
  - bound states  $\rightarrow$  negative phase shift
  - virtual states (**usually hidden !**)  $\rightarrow$  positive phase shift
  - resonances  $\rightarrow$  positive phase shift
  - left hand cut  $\rightarrow$  (empirically) negative phase shift (proved in quantum mechanical potential scatterings)

[T. Regge 1958 Nuovo Cimento]

# BRANCH CUT STRUCTURE OF PARTIAL WAVE $\pi N$ ELASTIC SCATTERING AMPLITUDE

[S. W. MacDowell 1959 PR][J. Kennedy and T. D. Spearman 1961 PR]



# TREE LEVEL LEFT-HAND CUT

- Tree level left-hand cut of  $S$ 
  - $(-\infty, (M_N - m_\pi)^2]$   $\rightarrow$  kinematic
  - $[(M_N^2 - m_\pi^2)^2/M_N^2, M_N^2 + 2m_\pi^2]$   $\rightarrow$   $u$  channel nucleon exchange  $\rightarrow$  very small
- The main contribution of  $f(s)$  (with a cut-off  $s_c$ )

$$f(s) = \frac{s}{\pi} \int_{s_c}^{(M_N - m_\pi)^2} \frac{\sigma(w)dw}{w(w-s)}$$

- The dispersion spectral function

$$\sigma(w) = \text{Im} \left\{ \frac{\ln |S_{\text{tree}}|}{2i\rho(w)} \right\} = -\frac{\ln |1 + 2i\rho(w)T_{\text{tree}}|}{2\rho(w)}$$

negative definite

- Right-hand inelastic cuts are omitted for the moment

# 3. Numerical results

# TREE-LEVEL QUALITATIVE ANALYSIS

- Values of the constants ( $s_c$  determined by  $N^*(1440)$  shadow pole position)

$$F = 0.0924 \text{ GeV}, g = 1.267, s_c = -0.08 \text{ GeV}^2$$

$$M_N = 0.9383 \text{ GeV}, m_\pi = 0.1396 \text{ GeV}$$

- $\mathcal{O}(p^2)$  K-Matrix phase shift:

$$T = T_{\text{tree}} / (1 - i\rho T_{\text{tree}}), \delta = \arctan(\rho T_{\text{tree}})$$

- Data from computer code SAID (WI 08)

<http://gwdac.phys.gwu.edu/>

- K-Matrix fit to  $S_{11}, S_{31}, P_{11}, P_{31}, P_{13}$  channels,  $W = \sqrt{s} \in [1.08, 1.16] \text{ GeV}$ ,  $\chi^2/\text{d.o.f.} = 1.850$ .

$$c_1 = -0.841, c_2 = 1.170, c_3 = -2.618, c_4 = 1.677$$





- Known poles [A.V. Anisovich et. al. 2012 Eur. Phys. J. A]

$$\sqrt{s_p}^{\text{II}} = M_p - i|\Gamma_{\pi N} - \Gamma_{\text{inel.}}|/2$$





# TREE LEVEL PHASE SHIFT RESULTS

$L_{2I} 2J$  convention,  $W = \sqrt{s}$ , data: green triangles [SAID: WI 08]



# TREE LEVEL PHASE SHIFT RESULTS

$L_{2I\ 2J}$  convention,  $W = \sqrt{s}$ , data: green triangles [SAID: WI 08]



# DISCREPANCIES IN $S_{11}$ AND $P_{11}$ CHANNELS

- Large missing positive contributions
- Possible interpretations
  - one loop contributions? numerical uncertainties?
  - contributions from other branch cuts?
  - hidden poles - virtual states, crazy resonances below threshold, or some extremely broad states?
- Once subtraction, logarithmic form  $\rightarrow$  **not sensitive to chiral orders and numerical details**



# RIGHT-HAND INELASTIC CUT

- Right-hand inelastic cut contribution  $\rightarrow$  positive definite

$$f_{R'}(s) = \frac{s}{\pi} \int_{(2m+M)^2}^{\Lambda_R^2} \frac{\sigma(w)dw}{w(w-s)}$$
$$\sigma(w) = - \left\{ \frac{\ln[\eta(w)]}{2\rho(w)} \right\}$$

- $\eta$ : inelasticity, from SAID WI 08 data and extrapolation
- Cut-off:  $\Lambda_R = 4.00\text{GeV}$

# RIGHT-HAND INELASTIC CUT




**Far from enough!!**

# 4. Hidden contributions

# FINDING $P_{11}$ HIDDEN POLE


- $P$ - wave:  $\delta(k) \sim \mathcal{O}(k^3)$
- Initially one resonance  $\rightarrow$  two virtual states  $\rightarrow$  one survives, the other is nearly absorbed by the point  $(M_N - m_\pi)^2$
- $s_c = -9 \text{ GeV}^2$ , virtual pole: 980 MeV,  $\chi_{P_{11}}^2/\text{d.o.f} = 0.201$ .
- An extra CDD pole is needed in  $P_{11}$  channel

[A. Gasparyan and M.F.M. Lutz 2010 NPA]



# $P_{11}$ CHANNEL: SHADOW POLE OF THE NUCLEON

- Analytical continuation:  $S^{\text{II}} = 1/S^{\text{I}}$ .  
Second sheet poles  $\rightarrow$  first sheet zeros.
- Expansion:  $S^{\text{I}} \sim a/(s - M_N^2) + b + \dots$
- Arbitrary non-zero  $b \rightarrow$  the virtual state
- Perturbative calculation  $\rightarrow$  virtual state at 976 MeV; fit  $\rightarrow$  980 MeV





# FINDING $S_{11}$ HIDDEN POLE

- $s_c = -0.08 \text{ GeV}^2$ ,  $\Lambda_R = 4.00 \text{ GeV}$ .
- Hidden pole  $\rightarrow$  a “crazy resonance” below threshold  
 $(0.861 \pm 0.053) - (0.130 \pm 0.075)i \text{ GeV}$

$s_c$ ( $\text{GeV}^2$ )	Pole position (GeV)	Fit quality $\chi^2/\text{d.o.f}$
-0.08	$0.808 - 0.055i$	0.109
-1	$0.822 - 0.139i$	0.076
-9	$0.883 - 0.195i$	0.034
$\infty$	$0.914 - 0.205i$	0.018

# $S_{11}$ CHANNEL: LOWEST POTENTIAL-NATURE RESONANCE?

- $S_{11}$  channel  $\rightarrow$  no  $s$ -channel intermediate states  $\rightarrow$  potential nature interaction
- Square-well potential ( $\mu$ : reduced mass)


$$U(r) = 2\mu V(r) = \begin{cases} -2\mu V_0 & (r \leq L), \\ 0 & (r > L), \end{cases}$$

- Phase shift ( $k' = (k^2 + 2\mu V_0)^{1/2}$ )

$$\delta_{\text{sw}}(k) = \arctan \left[ \frac{k \tan k'L - k' \tan kL}{k' + k \tan (kL) \tan (k'L)} \right]$$

# $S_{11}$ CHANNEL: LOWEST POTENTIAL-NATURE RESONANCE?

- Fit result (20 data):  $L = 0.829$  fm and  $V_0 = 144$  MeV,  $\chi_{\text{sw}}^2/\text{d.o.f} = 0.740$
- Pole position:  $k = -346i$  MeV  $\rightarrow 0.872 - 0.316i$  GeV.  
Hidden pole fit  $(0.861 \pm 0.053) - (0.130 \pm 0.075)i$  GeV



# 4. Summary

# SUMMARY

- PKU representation which separates phase shift contributions is employed to analyze  $\pi N$  elastic scatterings in  $s$  and  $p$  wave channels.
- The calculation of the left-hand cuts is under covariant baryon chiral perturbation theory at tree level.
- The  $S_{11}$  and  $P_{11}$  channels contain significant disagreements between “known poles + cut” and the experiment, missing large positive contributions. (reliable, independent of numerical details)
- $S_{11}$  channel contains a hidden resonance below threshold, while in  $P_{11}$  channel the nucleon pole induces a companionate virtual state.

**Thank you !!**

# Back up



- Intermediate particles

Channel	$I(J^P)$	Intermediate particles
$S_{11}$	$\frac{1}{2}(\frac{1}{2}^-)$	$N^*(1535), N^*(1650), N^*(1895)$
$S_{31}$	$\frac{3}{2}(\frac{1}{2}^-)$	$\Delta(1620), \Delta(1900)$
$P_{11}$	$\frac{1}{2}(\frac{1}{2}^+)$	$N, N^*(1440), N^*(1710), N^*(1880)$
$P_{31}$	$\frac{3}{2}(\frac{1}{2}^+)$	$\Delta(1910)$
$P_{13}$	$\frac{1}{2}(\frac{3}{2}^+)$	$N^*(1720), N^*(1900)$
$P_{33}$	$\frac{3}{2}(\frac{3}{2}^+)$	$\Delta(1232), \Delta(1600), \Delta(1920)$



# DETERMINATION OF COEFFICIENTS $c_i$ ?

- Set 1: this work
- Set 2:  $\mathcal{O}(p^3)$  fit in [Y. H. Chen et. al. 2013 PRD]
- Set 3: [D. Siemens et. al. 2017 PLB]
- Different choices have little impact on the left-hand cut contributions!



# $\mathcal{O}(p^3)$ PRELIMINARY RESULTS

- The same cut-off condition
- Chiral order does not impact on the existence of the  $S_{11}$  and  $P_{11}$  states
- $\mathcal{O}(p^3)$  greatly improves the fit quality in other channels that are impossible to fit the data at  $\mathcal{O}(p^2)$ , and there may be some indications of new hidden structures.

